

6197 Cornerstone Court, Ste 102 San Diego, CA 92121

Product Specifications

UltraPolymer Goat anti-Mouse IgG (H&L) – HRP

Description: UltraPolymer Goat anti-Mouse IgG (H&L) conjugated to HRP, affinity purified,

min x w/ bovine, horse, human, pig or rabbit serum proteins

2MH-050 Cat. No.: Amount: 50 mL Concentration: Ready to Use

Conjugate: **HRP**

Applications: IHC-P – optimized for use with autostainer or manual staining

Form: Liquid

Buffer: TBS. 1% (w/v) BSA. 0.1% Proclin 150. 10% heat-inactivated animal serum

Purification: Affinity purified using solid phase Mouse IgG

Affinity purified antibody is > 95% based on SDS-PAGE Purity:

Host: Goat Clonality: Polyclonal

Immunogen: Purified Mouse IgG, whole molecule

Storage:

Shelf-Life: 1 year from received date when stored at 2-8 °C.

Specificity: Based on IEP, this immunopolymer antibody reacts with heavy (y) chains on

mouse IgG, light chains on all mouse immunoglobulins

Cross-Reactivity: Based on IEP, no cross-reactivity is observed to non-immunoglobulin mouse

serum proteins, serum proteins from bovine, horse, human, pig/swine or

rabbit

Country of Origin: Serum was obtained from healthy animals of US origin and under the care of

a registered veterinarian.

Disclaimer: For In vitro Research Use Only. Not for diagnostic or therapeutic use. Suggested applications of


our products are not recommendations to use our products in violation of any patent or as a license under any patent of Cell IDx, inc. Product may not be resold or modified for resale without prior written approval of Cell IDx, inc. Liability of Cell IDx is limited to the purchase price

of the product.

Tonsil stained with mouse anti-human Ki-67 (clone B56, 0.4ug/ml) and detected with UltraPolymer goat anti-

mouse-HRP (RTU).

Staining on the BOND RX autostainer Antigen retrieval: ER2, 20 min Primary antibody: 30 min incubation Secondary antibody: 15 min incubation



Version 2.2: Revision date 2.5.21

www.cellidx.com info@cellidx.com 858-452-5800