



6197 Cornerstone Court, Ste 102  
San Diego, CA 92121


---

## Product Specifications

### UltraPolymer Donkey anti-Goat IgG (H&L) – HRP

Description:	UltraPolymer Donkey anti-Goat IgG (H&L) conjugated to HRP, affinity purified, min x w/ human, mouse, rabbit or rat IgG
Cat. No.:	2GH-050
Amount:	50 mL
Concentration:	Ready to Use
Conjugate:	HRP
Applications:	IHC-P – optimized for use with autostainer or manual staining
Form:	Liquid
Buffer:	TBS, 1% (w/v) BSA, 0.1% Proclin 150, 10% heat-inactivated animal serum
Purification:	Affinity purified using solid phase Goat IgG
Purity:	Affinity purified antibody is > 95% based on SDS-PAGE
Host:	Donkey
Clonality:	Polyclonal
Immunogen:	Purified Goat IgG, whole molecule
Storage:	2-8 °C
Shelf-Life:	1 year from received date when stored at 2-8 °C.
Specificity:	Based on IEP, this immunopolymer antibody reacts with heavy ( $\gamma$ ) chains on goat IgG, light chains on all goat immunoglobulins
Cross-Reactivity:	Based on IEP, no cross-reactivity is observed to non-immunoglobulin goat serum proteins, IgG from human, mouse, rabbit or rat
Country of Origin:	Serum was obtained from healthy animals of US origin and under the care of a registered veterinarian.
Disclaimer:	<i>For In vitro Research Use Only. Not for diagnostic or therapeutic use. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Cell IDx, inc. Product may not be resold or modified for resale without prior written approval of Cell IDx, inc. Liability of Cell IDx is limited to the purchase price of the product.</i>

Tonsil stained with goat anti-human IgG (1ug/ml) and detected with UltraPolymer donkey anti-goat-HRP (RTU).  
Staining on the BOND RX autostainer  
Antigen retrieval: ER2, 20 min  
Primary antibody: 30 min incubation  
Secondary antibody: 30 min incubation



Version 2.2: Revision date 2.5.21